

TRIVIAL MEASURES ARE NOT SO TRIVIAL

CHRISTOPHER P. PORTER

ABSTRACT. Although algorithmic randomness with respect to various non-uniform computable measures is well-studied, little attention has been paid to algorithmic randomness with respect to computable *trivial* measures, where a measure μ on 2^ω is trivial if the support of μ consists of a countable collection of sequences. In this article, it is shown that there is much more structure to trivial computable measures than has been previously suspected.

CONTENTS

1. Introduction	1
2. Computable Measures and Randomness	3
2.1. Computable measures on 2^ω	3
2.2. Notions of algorithmic randomness	4
2.3. Randomness and Turing functionals	7
3. Tally Functionals	7
4. Separating Randomness Notions via Trivial Measures	9
4.1. MLR and 2-randomness	9
4.2. MLR and weak-2 randomness	11
4.3. MLR and Schnorr randomness	12
5. Trivial Measures and Finite Distributive Lattices	13
References	23

1. INTRODUCTION

Algorithmic randomness with respect to non-uniform computable measures is well-studied. Although one can find definitions of biased random sequences in the work of von Mises, Ville, and Church, the first generally accepted definition of biased algorithmic randomness can be found in the final section of Martin-Löf’s groundbreaking 1966 paper, “The Definition of Random Sequences” ([ML66]), in which Martin-Löf considers definitions of algorithmic randomness for various Bernoulli measures. This study of definitions of biased randomness was continued in the 1970s by Levin and Zvonkin ([ZL70]) and Schnorr ([Sch71],[Sch77]). In particular, Levin, Zvonkin, and Schnorr studied measures that are *atomic*, i.e., measures μ for which there is some $A \in 2^\omega$ such that $\mu(\{A\}) > 0$, where A is called an *atom* of μ , denoted $A \in \text{Atoms}_\mu$.

Schnorr further studied what he called *discrete* measures, where μ is discrete if $\mu(\text{Atoms}_\mu) = 1$, or equivalently, if the support of μ is a countable collection of sequences. In [Sch77], Schnorr further claimed the following:

Claim. $\text{MLR}_\mu = \text{SR}_\mu$ if and only if μ is discrete.

Here MLR_μ denotes the collection of μ -Martin-Löf random sequences and SR_μ denotes the collection of μ -Schnorr random sequences.

Discrete measures were later referred to by Kautz's dissertation [Kau91] as *trivial*, the implication being that such measures have no interesting structural properties; once we've assigned all measure to some countable collection of points, there appears to be nothing left to say about the resulting measure.

The main goal of this paper is to show that this is *not* the case; there is much more structure to trivial measures than has been previously suspected. Not only do we show that the above claim of Schnorr's is false, but we also construct a number of trivial measures that allow us to separate notions of randomness. That is, we prove several theorems of the following form: given two randomness notions R^1 and R^2 such that $\text{R}_\mu^1 \subseteq \text{R}_\mu^2$ for every $\mu \in \mathcal{M}_c$, there is a trivial measure $\mu \in \mathcal{M}_c$ such that (i) $\text{R}_\mu^1 = \text{Atoms}_\mu$ and (ii) $\text{R}_\mu^2 \setminus \text{R}_\mu^1 \neq \emptyset$. In particular, we construct a trivial measure μ such that μ is constructed such that $\text{MLR}_\mu = \text{Atoms}_\mu$ and $\text{SR}_\mu \neq \text{Atoms}_\mu$. As further evidence of the non-triviality of trivial measures, we study a degree structure associated with a given trivial μ called the $LR(\mu)$ -degrees, showing that for each finite distributive lattice (L, \leq) , there is a computable trivial measure μ so that the collection of $LR(\mu)$ -degrees is isomorphic to (L, \leq) .

The outline of this paper is as follows. In Section 2, we provide the relevant technical background for the rest of the paper. Next, in Section 3, we discuss the main technique for constructing trivial measures that we employ throughout this paper, defining these measures in terms of what we call *tally functionals*. In Section 4, we separate various notions of randomness via trivial measures, including the separation of Martin-Löf randomness and Schnorr randomness that provides a counterexample to Schnorr's claim. Lastly, in Section 5, we discuss how to produce a trivial measure μ with an associated $LR(\mu)$ -degree structure that is isomorphic to a given finite distributive lattice.

We assume that the reader is familiar with the basics of computability theory: computable functions, partial computable functions, computably enumerable sets, Turing functionals, Turing degrees, the Turing jump, and so on (see, for instance, [Soa87]), as well as the basics of effective randomness (otherwise, we refer the reader to [DH10] or [Nie09]). We denote by 2^ω the set of infinite binary sequences, also known as Cantor space. We denote the set of finite strings by $2^{<\omega}$. \mathbb{Q}_2 is the set of dyadic rationals, i.e., multiples of a negative power of 2. Given $X \in 2^\omega$ and an integer n , $X \upharpoonright n$ is the string that consists of the first n bits of X , and $X(n)$ is the $(n+1)$ st of X (so that $X(0)$ is the first bit of X). If $\sigma, \tau \in 2^{<\omega}$, then $\sigma \preceq \tau$ means that σ is an initial segment of τ ; moreover, given $X \in 2^\omega$, $\sigma \prec X$ means that σ is an initial of X . Given a string σ , the basic open set determined by σ , is defined to be $[[\sigma]] = \{X : \sigma \prec X\}$. For $X, Y \in 2^\omega$, we define $X \oplus Y = \{2n : n \in X\} \cup \{2n+1 : n \in Y\}$. Given a collection $\{B_i\}_{i \in \omega} \subseteq 2^\omega$, $\bigoplus_{i \in \omega} B_i = \{\langle n, i \rangle : n \in B_i\}$, where $\langle \cdot, \cdot \rangle$ is some standard pairing function. For $A \in 2^\omega$, $A^{[i]} = \{n : \langle n, i \rangle \in A\}$, so that $A = \bigoplus_{i \in \omega} A^{[i]}$.

2. COMPUTABLE MEASURES AND RANDOMNESS

In this section, we review the relevant material on computability probability measures on 2^ω and the various notions of algorithmic randomness given in terms of these measures.

2.1. Computable measures on 2^ω . A probability measure on 2^ω assigns to each Borel subset of 2^ω a real in $[0, 1]$. It suffices to consider the restriction of probability measures to basic open subsets of 2^ω , for Caratheodory's theorem from classical measure theory ensures that a function μ defined on basic open sets and satisfying $\mu(\llbracket \sigma \rrbracket) = \mu(\llbracket \sigma 0 \rrbracket) + \mu(\llbracket \sigma 1 \rrbracket)$ for all $\sigma \in 2^{<\omega}$ can be uniquely extended to a probability measure. We can therefore represent measures as functions from strings to reals, where for all $\sigma \in 2^{<\omega}$, $\mu(\sigma)$ will denote the measure of $\llbracket \sigma \rrbracket$. This concise representation also allows us to talk about *computable* probability measures.

Definition 2.1. A probability measure μ on 2^ω is computable if $\sigma \mapsto \mu(\sigma)$ is computable as a real-valued function, i.e. if there is a computable function $\hat{\mu} : 2^{<\omega} \times \omega \rightarrow \mathbb{Q}_2$ such that

$$|\mu(\sigma) - \hat{\mu}(\sigma, i)| \leq 2^{-i}$$

for every $\sigma \in 2^{<\omega}$ and $i \in \omega$.

In what follows λ will refer exclusively to the Lebesgue measure on 2^ω , i.e., $\lambda(\sigma) = 2^{-|\sigma|}$ for each $\sigma \in 2^{<\omega}$. Moreover, \mathcal{M}_c denotes the collection of computable measures on 2^ω .

An important technique for defining a computable measure on 2^ω is to induce the measure by means of some Turing functional Φ . To do so, it must be the case that Φ is *almost total*, i.e., $\lambda(\text{dom}(\Phi)) = 1$. For our purposes, however, it suffices to restrict to truth-table functionals.

Definition 2.2. A Turing functional $\Phi : \subseteq 2^\omega \rightarrow 2^\omega$ is a truth-table functional (or *tt-functional*) if Φ is total.

Definition 2.3. Given a *tt-functional* $\Phi : 2^\omega \rightarrow 2^\omega$, the measure induced by Φ , denoted λ_Φ , is defined to be

$$\lambda_\Phi(\mathcal{X}) = \lambda(\Phi^{-1}(\mathcal{X}))$$

for every measurable $\mathcal{X} \subseteq 2^\omega$.

It is not hard to show that $\lambda_\Phi \in \mathcal{M}_c$ for each *tt-functional* Φ .

Apart from the Lebesgue measure, the computable measures that we will consider here are all atomic, and even trivial.

Definition 2.4. Let $\mu \in \mathcal{M}_c$.

- (i) μ is atomic if there is some sequence $A \in 2^\omega$ such that $\mu(\{A\}) > 0$.
- (ii) $A \in 2^\omega$ an atom of μ , denoted $A \in \text{Atoms}_\mu$, if $\mu(\{A\}) > 0$.
- (iii) μ is atomless if $\text{Atoms}_\mu = \emptyset$.
- (iv) μ is trivial if $\mu(\text{Atoms}_\mu) = 1$.

We will also refer to the atoms of μ as μ -atoms.

2.2. Notions of algorithmic randomness. Although there are many definitions of algorithmic randomness for elements of 2^ω , we restrict our attention to three such definitions: Martin-Löf randomness, Schnorr randomness, and weak 2-randomness.

Definition 2.5. Let $\mu \in \mathcal{M}_c$.

- (i) A μ -Martin-Löf test is a uniformly computable sequence $\{\mathcal{U}_i\}_{i \in \omega}$ of effectively open classes in 2^ω such that $\mu(\mathcal{U}_i) \leq 2^{-i}$ for every $i \in \omega$.
- (ii) $X \in 2^\omega$ is μ -Martin-Löf random if for every μ -Martin-Löf test $\{\mathcal{U}_i\}_{i \in \omega}$, we have $X \notin \bigcap_{i \in \omega} \mathcal{U}_i$.

The collection of μ -Martin-Löf random reals will be written as MLR_μ (we will simply write MLR when considering the Lebesgue measure). The following fact, the existence of a universal Martin-Löf test, is well-known and will prove to be useful here.

Proposition 2.6. For every $\mu \in \mathcal{M}_c$, there is a Martin-Löf test $\{\widehat{\mathcal{U}}_i\}_{i \in \omega}$ such that $x \in \text{MLR}_\mu$ if and only if $x \notin \bigcap_{i \in \omega} \widehat{\mathcal{U}}_i$.

We will make heavy use of the following lemma in Section 5:

Lemma 2.7. Given $\mu, \nu \in \mathcal{M}_c$, if we set $\rho = \frac{\mu + \nu}{2}$, then

$$\text{MLR}_\rho = \text{MLR}_\mu \cup \text{MLR}_\nu. \quad \text{¹}$$

Proof. If $X \notin \text{MLR}_\rho$, then $X \in \bigcap_{i \in \omega} \mathcal{U}_i$, where $\{\mathcal{U}_i\}_{i \in \omega}$ is the universal ρ -Martin-Löf test. But since $\rho(\mathcal{U}_i) \leq 2^{-i}$ for each $i \in \omega$, it follows that $\mu(\mathcal{U}_{i+1}) \leq 2^{-i}$ and $\nu(\mathcal{U}_{i+1}) \leq 2^{-i}$, and hence it follows that $X \notin \text{MLR}_\mu \cup \text{MLR}_\nu$.

Conversely, if $X \notin \text{MLR}_\mu \cup \text{MLR}_\nu$, then $X \in \bigcap_{i \in \omega} \mathcal{U}_i$ and $X \in \bigcap_{i \in \omega} \mathcal{V}_i$, where $\{\mathcal{U}_i\}_{i \in \omega}$ is the universal μ -Martin-Löf test and $\{\mathcal{V}_i\}_{i \in \omega}$ is the universal ν -Martin-Löf test. Then since $\mathcal{U}_i \cap \mathcal{V}_i \subseteq \mathcal{U}_i$ and $\mathcal{U}_i \cap \mathcal{V}_i \subseteq \mathcal{V}_i$, it follows that $\{\mathcal{U}_i \cap \mathcal{V}_i\}_{i \in \omega}$ is a ρ -Martin-Löf test, and hence $X \notin \text{MLR}_\rho$. \square

We will also draw heavily on the fact that there are Δ_2^0 Martin-Löf random sequences. Recall that a sequence $A \in 2^\omega$ is Δ_2^0 if there is a uniformly computable collection of finite sets $\{A_s\}_{s \in \omega}$ (called a Δ_2^0 approximation of A) such that

$$\lim_{n \rightarrow \infty} A_s(n) = A(n)$$

for every $n \in \omega$. For example, Chaitin's Ω , defined by

$$\Omega := \sum_{U(\sigma) \downarrow} 2^{-|\sigma|},$$

where U is a universal prefix-free Turing machine, is a Δ_2^0 Martin-Löf random sequence. In fact, $\Omega \equiv_T \emptyset'$.

The following results concerning μ -atoms and their relationship to μ -Martin-Löf randomness will be particularly useful.

Proposition 2.8. Let $\mu \in \mathcal{M}_c$ and $X \in 2^\omega$.

- (i) If $X \in \text{Atoms}_\mu$, then X is computable.

¹In general, we can consider any convex sum $\frac{\alpha\mu + (1-\alpha)\nu}{2}$ of μ and ν , as long as $\alpha \in [0, 1]$ is computable.

- (ii) If $X \in \text{Atoms}_\mu$, then $X \in \text{MLR}_\mu$.
- (iii) If X is computable and $\mu(\{X\}) = 0$, then $X \notin \text{MLR}_\mu$.

Part (ii) of the above proposition actually holds for all notions of randomness that we consider here, since for each such definition, the non-random sequences are captured in some set of μ -measure zero.

We will also consider relativized versions of Martin-Löf randomness. For $A \in 2^\omega$, a μ -Martin-Löf test relative to A is simply a uniformly A -computable sequence $\{\mathcal{U}_i^A\}_{i \in \omega}$ of $\Sigma_1^{0,A}$ classes in 2^ω such that $\mu(\mathcal{U}_i^A) \leq 2^{-i}$ for every $i \in \omega$. Moreover, $X \in 2^\omega$ is μ -Martin-Löf random relative to A , denoted $X \in \text{MLR}_\mu^A$, if $X \notin \bigcap_{i \in \omega} \mathcal{U}_i^A$ for any μ -Martin-Löf test relative to A $\{\mathcal{U}_i^A\}_{i \in \omega}$.

If we relativize Martin-Löf randomness to the halting set \emptyset' , then this gives rise to 2-randomness. We set $2\text{MLR} := \text{MLR}^{\emptyset'}$. It is immediate that $X \in 2\text{MLR}$ if and only if $X \in \text{MLR}^A$ for some $A \equiv_T \emptyset'$. It is not hard to show that $2\text{MLR} \subseteq \text{W2R}$.

One of the central results concerning relative randomness is known as “van Lambalgen’s theorem”.

Theorem 2.9 ([vL90]). *For every $A, B \in 2^\omega$,*

$$A \oplus B \in \text{MLR} \text{ if and only if } A \in \text{MLR}^B \ \& \ B \in \text{MLR}.$$

A related result is the following.

Theorem 2.10 ([Kau91]). *Given $A \in \text{MLR}$, then for each $i \in \omega$, $A^{[i]} \in \text{MLR}^{\oplus_{j \neq i} A^{[j]}}$. Further, for any finite $J \subseteq \omega$, $A^{[i]} \in \text{MLR}^{\oplus_{j \in J} A^{[j]}}$ for every $i \notin J$.*

Next we turn to Schnorr randomness, the definition of which is a slight variant of the definition of Martin-Löf randomness.

Definition 2.11. *Let $\mu \in \mathcal{M}_c$.*

- (i) *A μ -Schnorr test is a uniformly computable sequence $\{\mathcal{U}_i\}_{i \in \omega}$ of effectively open classes in 2^ω such that $\mu(\mathcal{U}_i) = 2^{-i}$ for every $i \in \omega$.*
- (ii) *$X \in 2^\omega$ is μ -Schnorr random if for every μ -Schnorr test $\{\mathcal{U}_i\}_{i \in \omega}$, we have $X \notin \bigcap_{i \in \omega} \mathcal{U}_i$.*

Let SR_μ be the class of μ -Schnorr random sequences. One can readily observe that $\text{MLR}_\mu \subseteq \text{SR}_\mu$ for each $\mu \in \mathcal{M}_c$. In order to separate Martin-Löf randomness and Schnorr randomness with respect to a trivial computable measure μ , we need the following result.

Proposition 2.12. [NST05] *Given $\mu \in \mathcal{M}_c$, every $X \in \text{SR}_\mu \setminus \text{MLR}_\mu$ is high, i.e. X computes a function that dominates all computable functions.*

We will review the proof of this proposition, as the details will be useful when we provide a counterexample to Schnorr’s claim in Section 4.

Proof of Proposition 2.12. Let $X \in \text{SR}_\mu \setminus \text{MLR}_\mu$, and let $\{\mathcal{U}_i\}_{i \in \omega}$ be a universal μ -Martin-Löf test. Then we define $f \leq_T X$ as follows:

$$f(n) = \text{the least } s \text{ such that } (\exists k)[X \upharpoonright k] \subseteq \mathcal{U}_{n,s}.$$

We claim that f dominates all computable functions. Suppose, for the sake of contradiction, that there is some computable function g such that $f(n) \leq g(n)$ for infinitely many n . Then $\{\mathcal{U}_{n,g(n)}\}_{n \in \omega}$ is in fact a Schnorr test, and moreover, there are infinitely many n such that $X \in \mathcal{U}_{n,g(n)}$, which is, in fact, sufficient to show that X is not Schnorr random (for instance, see [DH10], Theorem 7.1.10). Thus, X computes a function that dominates all computable functions, and hence X is high. \square

The third definition of randomness that we will consider here is weak 2-randomness.

Definition 2.13. Let $\mu \in \mathcal{M}_c$.

- (i) A generalized μ -Martin-Löf test is a uniformly computable sequence $\{\mathcal{U}_i\}_{i \in \omega}$ of effectively open classes in 2^ω such that $\lim_{i \rightarrow \infty} \mu(\mathcal{U}_i) = 0$.
- (ii) $X \in 2^\omega$ is μ -weakly 2-random if for every generalized μ -Martin-Löf test $\{\mathcal{U}_i\}_{i \in \omega}$, we have $X \notin \bigcap_{i \in \omega} \mathcal{U}_i$.

Let W2R_μ denote the collection of μ -weakly 2-random sequences. Note that every generalized μ -Martin-Löf test $\{\mathcal{U}_i\}_{i \in \omega}$ yields a Π_2^0 class of μ -measure zero, namely $\bigcap_i \mathcal{U}_i$, and conversely, every Π_2^0 class of μ -measure zero can be obtained in this way by a generalized μ -Martin-Löf test. It is immediate that $\text{W2R}_\mu \subseteq \text{MLR}_\mu$, since the collection of sequences captured by a μ -Martin-Löf defines a Π_2^0 class of μ -measure zero. We now consider the key result concerning the relationship between W2R_μ and MLR_μ .

Definition 2.14. $X, Y \in 2^\omega$ form a minimal pair in the Turing degrees if $A <_T X$ and $A <_T Y$ implies that $A \equiv_T \emptyset$.

Theorem 2.15 (Downey, Nies, Weber, Yu [DNWY06]; Hirschfeldt, Miller (unpublished)). For $\mu \in \mathcal{M}_c$, if X is not computable, then $X \in \text{W2R}_\mu$ if and only if $X \in \text{MLR}_\mu$ and X and \emptyset' form a minimal pair.

The proof of this result as found in the literature is only for the case of the Lebesgue measure, but it's not difficult to extend to any $\mu \in \mathcal{M}_c$. First, we follow the original proof, with several modifications.

Theorem 2.16 (Downey, Nies, Weber, and Yu [DNWY06]). For $\mu \in \mathcal{M}_c$, if $X \in \text{W2R}_\mu$ and X is not computable, then X and \emptyset' form a minimal pair.

Proof Sketch. We modify the proof given by Downey, Nies, Weber, and Yu for the case that $\mu = \lambda$. If $A \in 2^\omega$ is Δ_2^0 , $Z \in \text{W2R}_\mu$, and $\Phi^Z = A$ for some Turing functional A , then we argue that A is computable. Towards this end, we define

$$\mathcal{S} = \{X : \forall n \forall s \exists t > s (\Phi^X(n)[t] \downarrow = A_t(n))\},$$

which is Π_2^0 and contains Z . Since $Z \in \text{W2R}_\mu$, it follows that $\mu(\mathcal{S}) > 0$. Now the only difference between the original proof and the situation here is that μ may be atomic, so that there is some μ -atom $Y \in \mathcal{S}$. But in this case we are done: since μ is computable, it follows that Y is computable. Then $\Phi^Y = A$, and hence A is computable.

In the case that \mathcal{S} contains no atoms, the proof proceeds exactly as in the case of the Lebesgue measure: by a ‘‘majority vote’’ argument, which shows that that one can compute values of A using the majority of sequences in a set of positive measure, one shows that A is computable. See, for instance, the proof of Theorem 7.2.8. in [DH10] for the details. \square

Theorem 2.17. [Hirschfeldt, Miller (unpublished)] Let $\mu \in \mathcal{M}_c$. For any Σ_3^0 class $\mathcal{S} \subseteq 2^\omega$ such that $\mu(\mathcal{S}) = 0$, there is a noncomputable c.e. set A such that $A \leq_T X$ for every non-computable $X \in \text{MLR}_\mu \cap \mathcal{S}$.

Proof Sketch. The proof proceeds exactly in the case as the case of the Lebesgue measure, since $X \in \text{MLR}_\mu \cap \mathcal{S}$ implies that $\mu(\{X\}) = 0$ and hence X is not a μ -atom. See the proof of Theorem 7.2.11 of [DH10]. \square

Proof of Theorem 2.15. (\Rightarrow) This is simply Theorem 2.16.

(\Leftarrow) Suppose that $X \in \text{MLR}_\mu \setminus \text{W2R}_\mu$. Then there is a Π_2^0 μ -null set \mathcal{S} such that $X \in \mathcal{S}$, and so by Theorem 2.17, there is some noncomputable c.e. set A such that $A \leq_T X$. Therefore, X and \emptyset' do not form a minimal pair. \square

2.3. Randomness and Turing functionals. Many of the results in the sequel depend crucially on the following preservation of randomness theorems.

Theorem 2.18 (Preservation of Martin-Löf Randomness). *If Φ is a tt-functional, then $X \in \text{MLR}$ implies $\Phi(X) \in \text{MLR}_{\lambda_\Phi}$.*

Theorem 2.19 (Preservation of Schnorr Randomness). *If Φ is an tt-functional, then $X \in \text{SR}$ implies $\Phi(X) \in \text{SR}_{\lambda_\Phi}$.*

For the proofs of these results, see, for instance [BP12].

We will also make use of a relative version of the preservation of Martin-Löf randomness: For any $A \in 2^\omega$, if Φ is a tt-functional and $X \in \text{MLR}^A$, then $\Phi(X) \in \text{MLR}_{\lambda_\Phi}^A$. A partial converse to the preservation of Martin-Löf randomness is the following.

Theorem 2.20. *For Φ an almost total functional and $Y \in 2^\omega$, if $Y \in \text{MLR}_{\lambda_\Phi}$, then there is some $X \in \text{MLR}$ such that $\Phi(X) = Y$.*

A relativized version of Theorem 2.20 also holds, and combining this result with the relativized version of the preservation of Martin-Löf randomness, one can prove the following:

Theorem 2.21. *Given $X, A \in 2^\omega$ and a tt-functional Φ , if $\Phi(X)$ is not computable and $\Phi^{-1}(\Phi(X)) = \{X\}$, then $X \in \text{MLR}^A$ if and only if $\Phi(X) \in \text{MLR}_{\lambda_\Phi}^A$.*

3. TALLY FUNCTIONALS

The main tool used in the construction of various trivial measures are what we refer to here as “tally functionals”. Given a Δ_1^0 formula $\Theta(X, y, z)$ with free first-order variables y and z and free second-order variable X , we define an auxiliary function $\theta(A, n) : 2^\omega \times \omega \rightarrow \omega$ by

$$\theta(A, n) = \begin{cases} \text{the least } s \text{ such that } \Theta(A, n, s) & \text{if } s \text{ exists} \\ +\infty & \text{otherwise} \end{cases}$$

Then the tally functional Φ_Θ determined by the formula Θ is defined to be

$$\Phi_\Theta(A) = 1^{\theta(A,0)} 0 1^{\theta(A,1)} 0 1^{\theta(A,2)} 0 \dots,$$

where

$$\Phi_\Theta(A) = 1^{\theta(A,0)} 0 1^{\theta(A,1)} 0 \dots 1^{\theta(A,k)} 0 1^\omega$$

if $\theta(A, k+1) = +\infty$ (and is the least n such that $\theta(A, n) = +\infty$).

A useful example that we will use repeatedly in Section 5 is a tally functional defined in terms of the approximation of a Δ_2^0 sequence $A \in \text{MLR}$. Let $\{A_s\}_{s \in \omega}$ be a Δ_2^0 approximation of A . Without loss of generality, we can assume that $A_s \neq A_{s+1}$ for every s . If we define the formula $\Theta(A, n, s)$ so that

$$\Theta(A, n, s) \text{ holds if and only if } A \upharpoonright n = A_s \upharpoonright n,$$

then $\theta(A, n)$ is the first stage s such that $A \upharpoonright n = A_s \upharpoonright n$. Let Φ_A be the resulting tally functional.

Lemma 3.1. *Suppose that $A \in \Delta_2^0 \cap \text{MLR}$, and let Φ_A be the tally functional defined above.*

- (i) *If $X = A_s$ for some s , then there is some m such that for every $n \geq m$, $\theta(X, n) = \theta(X, m)$, i.e. the function $f(x) = \theta(X, x)$ is eventually constant.*
- (ii) *If $X \neq A$ and $X \neq A_s$ for each s , then $\theta(X, n) = +\infty$ for some n .*
- (iii) *The function $g(x) = \theta(A, x)$ is not computable, nor is it dominated by any computable function.*

Proof. (i) Let s be least such that $X = A_s$. If $s = 0$, then since for all n , $X \upharpoonright n = A_0 \upharpoonright n$, it follows that $\theta(X, n) = 0$ for all n . Now suppose that $s > 0$. Then there is some m such that $X \upharpoonright m \neq A_{s-1} \upharpoonright m$. It thus follows that $X \upharpoonright n = A_s \upharpoonright n$ for all $n \geq m$, which implies that $\theta(X, n) = s$ for all $n \geq m$.

(ii) First we establish the following claim: There is some k such that for every s , $X \upharpoonright k \neq A_s \upharpoonright k$. Suppose not, so that for every k , there is some s such that $X \upharpoonright k = A_s \upharpoonright k$. But this implies that for every k , there exist infinitely many s such that $X \upharpoonright k = A_s \upharpoonright k$, since (a) $X \neq A_s$ for every s and (b) $X \upharpoonright k = A_s \upharpoonright k$ implies that $X \upharpoonright j = A_s \upharpoonright j$ for every $j < k$. From this it follows that the A_s 's (viewed as rational numbers) converge to X (viewed as a real number). But the A_s 's also converge to A , and thus it follows that $X = A$, contradicting our hypothesis. Now let k be least such $X \upharpoonright k \neq A_s \upharpoonright k$ for every s . Then it follows that $\theta(X, k) = +\infty$.

(iii) If $g(n) = \theta(A, n)$ were computable, then since

$$A_{g(n)} \upharpoonright n = A_{\theta(A, n)} \upharpoonright n = A \upharpoonright n,$$

this would imply that A is computable, contradicting our assumption that $A \in \text{MLR}$. The proof that g is not dominated by any computable function is just given by the standard proof used to show that every non-computable Δ_2^0 sequence is hyperimmune. See, for instance, [Nie09] Theorem 1.5.12. □

Theorem 3.2. (i) *The measure μ induced by Φ_A is trivial.*

(ii) *If $A \in \text{MLR}$, then $\text{MLR}_\mu \setminus \text{Atoms}_\mu = \{\Phi_A(A)\}$.*

Proof. (i) Given input X , there are three cases to consider to determine the output $\Phi_\Theta(X)$:

Case 1: $X = A_s$ for some s . In this case, by Lemma 3.1 (i), the function $f(x) = \theta(X, x)$ is eventually constant, and thus $\Phi_\Theta(X) = \sigma 1^k 0 1^k 0 1^k 0 \dots$ for some $\sigma \in 2^{<\omega}$, which is clearly computable.

Case 2: $X \neq A$ and $X \neq A_s$ for every s . Then by Lemma 3.1 (ii), there is some n such that $\theta(X, n) = +\infty$, and hence $\Phi_\Theta(A) = \sigma 1^\omega$ for some $\sigma \in 2^{<\omega}$, which is computable.

Case 3: $X = A$. By Lemma 3.1 (iii), since the function $g(n) = \theta(A, n)$ is not computable, it follows that

$$\Phi_\Theta(A) = 1^{\theta(A,0)} 0 1^{\theta(A,1)} 0 1^{\theta(A,2)} 0 \dots,$$

is not a computable sequence.

For every $B \in \text{MLR}$ such that $B \neq A$, we must be in Case 2, as $B \neq A_s$ for every s . Thus $\Phi_\Theta(B) = \sigma 1^\omega$ for some $\sigma \in 2^{<\omega}$. Setting

$$\mathcal{S} = \{Y : (\exists \sigma \in 2^{<\omega})[Y = \sigma 1^\omega]\},$$

we have

$$\text{MLR} \setminus \{A\} \subseteq \Phi_\Theta^{-1}(\mathcal{S}),$$

from which it follows that

$$1 = \lambda(\text{MLR} \setminus \{A\}) \leq \lambda(\Phi_\Theta^{-1}(\mathcal{S})).$$

Since $\mu = \lambda_{\Phi_\Theta}$ assigns measure one to the countable collection \mathcal{S} , it follows that μ is trivial.

(ii) First, $\Phi_\Theta(A) \in \text{MLR}_\mu$ by the preservation of Martin-Löf randomness. In addition, $\Phi_\Theta(A) \notin \text{Atoms}_\mu$, for otherwise as $\Phi_\Theta(A)$ would be computable by Proposition 2.8 (i), and hence the function $g(n) = \theta(A, n)$ would be computable, contradicting Lemma 3.1 (iii).

Next, if $X \in \text{MLR}_\mu \setminus \text{Atoms}_\mu$, then by Theorem 2.20, $X \in \text{MLR}_\mu$ implies that $X = \Phi_\Theta(Y)$ for some $Y \in \text{MLR}$. But since $X \notin \text{Atoms}_\mu$, X is not computable. In particular, X does not have the form $\sigma 1^\omega$ for any $\sigma \in 2^{<\omega}$. It follows that Y cannot fall under Case 2, and since no $Y \in \text{MLR}$ falls under Case 1, it must be that $Y = A$. Thus, $X = \Phi_\Theta(A)$. □

We will revisit this result in Section 5, when we consider the LR-degree structures associated to different trivial measures. For other applications of tally functionals, see [BP12].

4. SEPARATING RANDOMNESS NOTIONS VIA TRIVIAL MEASURES

In this section, we prove three results of the following form: Let \mathbf{R}^1 and \mathbf{R}^2 be two notions of randomness such that $\mathbf{R}_\mu^1 \subseteq \mathbf{R}_\mu^2$ for every $\mu \in \mathcal{M}_c$. Then there is a measure $\mu \in \mathcal{M}_c$ such that (i) $\mathbf{R}_\mu^1 = \text{Atoms}_\mu$ and (ii) $\mathbf{R}_\mu^2 \setminus \mathbf{R}_\mu^1 \neq \emptyset$. Moreover, we will be able to conclude that μ is trivial, since by condition (i), $\mu(\text{Atoms}_\mu) = 1$.

4.1. MLR and 2-randomness. In order to separate MLR and 2MLR via a trivial measure, we prove a more general result by modifying the construction of a trivial measure in the previous section, and then we apply van Lambalgen's Theorem.

Theorem 4.1. *Let $A \in \text{MLR} \cap \Delta_2^0$. Then there is a trivial measure $\mu \in \mathcal{M}_c$ such that*

- (i) $\text{MLR}_\mu^A = \text{Atoms}_\mu$, and
- (ii) $\text{MLR}_\mu^A \setminus \text{MLR}_\mu^A \neq \emptyset$.

Proof. We define a new functional Ψ that on input $X \oplus Y$ behaves much like the tally functional Φ_A defined in Section 3. However, instead having our functional output a sequence of blocks 1s separated by individual 0s, Ψ will output blocks consisting the bits of Y . Specifically, suppose that

$$\Phi_A(X) = 1^{t_0}01^{t_1}01^{t_2}0 \dots 1^{t_i}0 \dots$$

Then we have

$$\Psi(X \oplus Y) = y_0^{t_0} y_1^{t_1} y_2^{t_2} \dots y_i^{t_i} \dots$$

where $y_i = Y(i)$ for every i . Note that Ψ is total, since Φ_A is total. Let μ be the induced measure λ_Ψ . Given input $X \oplus Y$, there are three relevant cases to consider, corresponding to the three cases we considered in the proof of Theorem 3.2.

Case 1: $X = A_s$ for some s . Then for any $Y \in 2^\omega$, there is some k such that

$$\Psi(A_s \oplus Y) = y_0^{t_0} y_1^{t_1} y_2^{t_2} \dots y_i^k y_{i+1}^k y_{i+2}^k \dots$$

That is, the lengths of the blocks of the y_i 's eventually stabilize (just as the function $\theta(A_s, n)$ in Case 1 of the proof of Theorem 3.2 eventually stabilizes).

Case 2: $X \neq A$ and $X \neq A_s$ for every s . Then there is some $\sigma \in 2^{<\omega}$ and $i \in \omega$ such that after some stage, Ψ will output the same bit y_i forever, i.e.,

$$\Psi(X \oplus Y) = \sigma(y_i)^\omega.$$

Case 3: $X = A$. Then using the same function $g(n) = \theta(A, n)$ from Lemma 3.1 (iii), we have

$$\Psi(A \oplus Y) = y_1^{\theta(A,0)} y_2^{\theta(A,1)} y_3^{\theta(A,2)} \dots$$

Note further that if Y has only finitely many 0s or finitely many 1s, then $\Psi(A \oplus Y)$ will eventually stabilize. In the case that Y has infinitely 0s and 1s, then there is some function $f : \omega \rightarrow \omega$ such that

$$\Psi(A \oplus Y) = b_0^{f(0)} b_1^{f(1)} b_2^{f(2)} \dots y_i^{f(i)} \dots$$

where $b_i \neq b_{i+1}$ for each $i \in \omega$. It is not hard to see that $\theta(A, n) \leq f(n)$ for every $n \in \omega$, from which it follows that f is not computable by Lemma 3.1 (iii), and thus $\Psi(A \oplus Y)$ is not computable.

Now we verify (i) and (ii). For (i), it is clear that $\text{Atoms}_\mu \subseteq \text{MLR}_\mu^A$. Given $Z \in \text{MLR}_\mu^A$, by Theorem 2.20 relativized to A , there must be some A -random sequence $X \oplus Y$ such that $\Psi(X \oplus Y) = Z$. If Z is not computable, it must be the case that either $X = A$ or $X = A_s$ for some $s \in \omega$, for otherwise we would be in Case 2, so that $\Psi(X \oplus Y) = Z$ would be computable. But $A \oplus Y$ is not Martin-Löf random relative to A , nor is $A_s \oplus Y$ for any $s \in \omega$. Thus, Z must be computable, and so by Proposition 2.8 (iii), $Z \in \text{Atoms}_\mu$.

For (ii), let $B \in \text{MLR}^A$. Then $A \oplus B \in \text{MLR}$ by van Lambalgen's Theorem (Theorem 2.9). It follows from the preservation of Martin-Löf randomness that $\Psi(A \oplus B) \in \text{MLR}_\mu$. Since B has infinitely many 0s and 1s, by the discussion in Case 3, $\Psi(A \oplus B)$ is not computable. By Proposition 2.8 (i), $\Psi(A \oplus B) \notin \text{Atoms}_\mu$, and so by part (i), $\Psi(A \oplus B) \notin \text{MLR}_\mu^A$. \square

We now have the following corollary:

Corollary 4.2. *There is a trivial measure μ such that*

- (i) $2\text{MLR}_\mu = \text{Atoms}_\mu$, and
- (ii) $\text{MLR}_\mu \setminus 2\text{MLR}_\mu \neq \emptyset$.

Proof. Choose any $A \in \Delta_2^0 \cap \text{MLR}$ such that $A \equiv_T \emptyset'$ (for instance, let $A = \Omega$) and apply Theorem 4.1. \square

4.2. MLR and weak-2 randomness. We can improve Corollary 4.2 by replacing 2MLR with W2R. However, we need to use a different technique to do so: we will use the characterization provided by Theorem 2.15, that for each non-computable X , $X \in \text{W2R}_\mu$ if and only $X \in \text{MLR}_\mu$ and X forms a minimal pair with \emptyset' .

Theorem 4.3. *There is a trivial measure $\mu \in \mathcal{M}_c$ such that*

- (i) $\text{W2R}_\mu = \text{Atoms}_\mu$, and
- (ii) $\text{MLR}_\mu \setminus \text{W2R}_\mu \neq \emptyset$.

Proof. Let $A \in \text{MLR} \setminus \text{W2R}$. Then by Theorem 2.15 there is some Turing functional Γ and a non-computable Δ_2^0 sequence B such that $\Gamma(A) = B$. Let $\Theta(X, n, s)$ be such that

$$\Theta(X, n, s) \text{ holds if and only if } (\forall k < n) \Gamma_s(X)(k) \downarrow = B_s(k)$$

where $\{B_s\}_{s \in \omega}$ is some fixed Δ_2^0 approximation of B , where without loss of generality $B_s \neq B_{s+1}$ for every $s \in \omega$. Then it follows that θ is defined to be

$$\theta(X, n) = \begin{cases} \text{the least } s \text{ such that } (\forall k < n) \Gamma_s(X)(k) \downarrow = B_s(k) & \text{if } s \text{ exists} \\ +\infty & \text{otherwise} \end{cases}$$

Given $X \in 2^\omega$, to compute $\Phi_\Theta(X)$, we have four cases to consider, each depending on the behavior of Γ on input X .

Case 1: $\Gamma(X) \uparrow$. Then there is some least n such that $\Gamma(X)(n) \uparrow$, and so $\theta(X, k) = +\infty$ for some $k \leq n+1$ (there may be some $k < n+1$ such that $\Gamma(X)(k) \downarrow$ but $\Gamma(X)(k) \neq B_s(k)$ for every $s \in \omega$). Consequently,

$$(1) \quad \Phi_\Theta(X) = \sigma 1^\omega$$

for some $\sigma \in 2^{<\omega}$.

Case 2: $\Gamma(X) \downarrow = B_t$ for some $t \in \omega$. First observe that for any $k > t$, $\Gamma(X)(k)$ halts in at least k steps, given the standard convention on the running times of computations (i.e. the running time must always exceed the size of the input and the output). Since $B_s \neq B_{s+1}$ for every $s \in \omega$, there is some least n such that for all sufficiently large $s \geq n$

$$(\forall k < n) [\Gamma_s(k) \downarrow = B_t(k)]$$

but

$$(\exists j < n) [B_s(j) \neq B_t(j)].$$

That is, the functional Φ_Θ on input X waits to see $\Gamma_s(X)$ agree with B_s , but for sufficiently large stages, $\Gamma_s(X)$ will only agree with B_t . Hence for the n given above, we have $\theta(X, n) = +\infty$, and thus (1) holds.

Case 3: $\Gamma(X) \downarrow$ but $\Gamma(X) \neq B$ and $\Gamma(X) \neq B_s$ for every $s \in \omega$. In this case we have

$$(2) \quad (\exists n)(\forall s)(\exists k < n) [\Gamma_s(X)(k) \neq B_s(k)].$$

The argument to establish this claim is the same as the one we gave in the proof of Lemma 3.1 (ii). Let n be the least number satisfying (2). Then we have $\theta(X, n) = +\infty$, and thus (1) holds.

Case 4: $\Gamma(X)\downarrow = B$. Setting $h(n) := \theta(X, n)$, we have

$$\Phi_{\Theta}(X) = 1^{h(0)}01^{h(1)}0\dots,$$

so that $\Phi_{\Theta}(X) \geq_T h$. But for every $n \in \omega$,

$$B_{h(n)}\upharpoonright n = \Gamma_{h(n)}(X)\upharpoonright n = \Gamma(X)\upharpoonright n = B\upharpoonright n,$$

so it follows that $\Phi_{\Theta}(X) \geq_T B$.

To establish (i), $\text{Atoms}_{\mu} \subseteq \text{W2R}_{\mu}$ is immediate. For the other direction, consider $Y \in \text{W2R}_{\mu}$, which by Theorem 2.15 forms a minimal pair with \emptyset' . Since $Y \in \text{MLR}_{\mu}$, by Theorem 2.20 there is some $X \in \text{MLR}$ such that $\Phi_{\Theta}(X) = Y$.

Applying the functional Γ to X yields one of two general outcomes: either one of Case 1, 2, or 3 occurs, in which case

$$Y = \Phi_{\Theta}(X) = \sigma 1^{\omega}$$

and hence $Y \in \text{Atoms}_{\mu}$ by Proposition 2.8 (ii), or we are in Case 4. But in this case, $Y = \Phi_{\Theta}(X) \geq_T B$, contradicting the fact that Y forms a minimal pair with \emptyset' . So we must have $Y \in \text{Atoms}_{\mu}$.

For part (ii), if we take the original $A \in \text{MLR} \setminus \text{W2R}$ such that $\Gamma(A) = B$ that we started with, by the preservation of Martin-Löf randomness, $\Phi_{\Theta}(A) \in \text{MLR}_{\mu}$. Further, by Case 4 above, $\Phi_{\Theta}(A) \geq_T B$, so $\Phi_{\Theta}(A)$ does not form a minimal pair with \emptyset' . Hence $\Phi_{\Theta}(A) \notin \text{W2R}_{\mu}$. \square

4.3. MLR and Schnorr randomness. We end this section by using a tally functional to show that Schnorr's claim, namely that $\text{MLR}_{\mu} = \text{SR}_{\mu}$ for a computable measure μ if and only if μ is trivial, is false. Here we will use yet another technique for constructing the requisite tally functional.

Theorem 4.4. *There is a trivial measure $\mu \in \mathcal{M}_c$ such that*

- (i) $\text{MLR}_{\mu} = \text{Atoms}_{\mu}$, and
- (ii) $\text{SR}_{\mu} \setminus \text{MLR}_{\mu} \neq \emptyset$.

Proof. To construct the desired measure μ , we define a tally functional in terms of a universal μ -Martin-Löf test $\{\mathcal{U}_i\}_{i \in \omega}$. Let $\Theta(A, n, s)$ be such that

$$\Theta(A, n, s) \text{ if and only if } (\exists k < s)[[A\upharpoonright k] \subseteq \mathcal{U}_{n,s}].$$

As above, $\theta(A, n)$ is the least such s such that $\Theta(A, n, s)$ holds, or is $+\infty$ if it fails to hold for every s .

There are two cases of interest to us (the case that $X \notin \text{MLR} \cup \text{SR}$ has no bearing on the result here).

Case 1: $X \in \text{MLR}$. In this case, there is some least n such that $X \notin \mathcal{U}_n$, and hence $\theta(X, n) = +\infty$, so that

$$\Phi_{\Theta}(X) = \sigma 1^{\omega}$$

for some $\sigma \in 2^{<\omega}$.

Case 2: $X \in \text{SR} \setminus \text{MLR}$. Then $X \in \bigcap_{i \in \omega} \mathcal{U}_i$, and moreover, by the proof of Theorem 2.12, the function $f \leq_T X$ such that

$$f(n) = \text{the least } s \text{ such that } (\exists k)[X \upharpoonright k] \subseteq \mathcal{U}_{n,s}.$$

dominates all computable functions. It follows that $\theta(X, n) = f(n)$, so that

$$\Phi_\Theta(X) = 1^{f(0)} 0 1^{f(1)} 0 \dots$$

is not computable.

To verify (i), as above $\text{Atoms}_\mu \subseteq \text{MLR}_\mu$ is immediate. Now given $Z \in \text{MLR}_\mu$, by Theorem 2.20, there is some $X \in \text{MLR}$ such that $\Phi_\Theta(X) = Z$. But we are in Case 1, so $Z = \sigma 1^\omega$ for some $\sigma \in 2^{<\omega}$. By Proposition 2.8 (iii), $Z \in \text{Atoms}_\mu$.

For (ii), given any $X \in \text{SR} \setminus \text{MLR}$, by the preservation of Schnorr randomness, we have $\Phi_\Theta(X) \in \text{SR}_\mu$. Further, since we are in Case 2, $\Phi_\Theta(X)$ is not computable, and thus by Proposition 2.8 (i) $\Phi_\Theta(X) \notin \text{Atoms}_\mu$. Thus by part (i), $\Phi_\Theta(X) \notin \text{MLR}_\mu$. □

5. TRIVIAL MEASURES AND FINITE DISTRIBUTIVE LATTICES

As further evidence of the non-triviality of trivial measures, we show that a trivial measure gives rise to a certain structure, which varies as we consider different trivial measures. Specifically, if one considers the *LR*-degrees (or “low-for-random” degrees) associated with MLR_μ for a class of trivial measures μ , one finds that different trivial measures can give rise to non-isomorphic *LR*-degree structures.

Nies [Nie05] gave the following definition in the context of Martin-Löf randomness with respect to the Lebesgue measure. We say that A is *LR-reducible* to B , denoted $A \leq_{LR} B$ if

$$\text{MLR}^B \subseteq \text{MLR}^A.$$

The intuitive idea is that B is more powerful than A as an oracle, as B de-randomizes more sequences than A does. Further, we say that A is *LR-equivalent* to B , denoted $A \equiv_{LR} B$ if and only if $A \leq_{LR} B$ and $B \leq_{LR} A$. The collection of *LR*-equivalence classes is called the collection of *LR-degrees*, denoted by \mathcal{D}_{LR} . Like the Turing degrees \mathcal{D}_T , the *LR*-degrees form an uncountable upper semilattice. However, unlike the structure of the Turing degrees (\mathcal{D}_T, \leq_T) , the structure of $(\mathcal{D}_{LR}, \leq_{LR})$ is not well-understood.

We can extend the definition of \leq_{LR} to Martin-Löf randomness with respect to any $\mu \in \mathcal{M}_c$ as follows. For $\mu \in \mathcal{M}_c$ and $A, B \in 2^\omega$, we say that A is *LR(μ)-reducible* to B , denoted $A \leq_{LR(\mu)} B$ if

$$\text{MLR}_\mu^B \subseteq \text{MLR}_\mu^A.$$

Similarly, we can define the *LR(μ)-degrees*, denoted $\mathcal{D}_{LR(\mu)}$, just as we defined the *LR*-degrees above. Interestingly, for certain choices of $\mu \in \mathcal{M}_c$, the structure $(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)})$ is very simple. Let’s consider some examples.

Example 1. Let $\mu \in \mathcal{M}_c$ be such that $\mu(\text{Atoms}_\mu) = 1$ and $\text{Atoms}_\mu = \text{MLR}_\mu$. Then $\mathcal{D}_{LR(\mu)}$ consists of a single equivalence class, consisting of all of 2^ω . The reason is that if $\mu(\{X\}) > 0$, then $X \in \text{MLR}_\mu^A$ for every $A \in 2^\omega$.

Example 2. If μ is the measure induced by the tally functional Φ_A for $A \in \Delta_2^0 \cap \text{MLR}$ (as in the example from Section 3), then $\mathcal{D}_{LR(\mu)}$ consists of exactly two elements. If we set $A^* := \Phi_A(A)$, then by Theorem 3.2 (ii),

$$\text{MLR}_\mu = \{A^*\} \cup \text{Atoms}_\mu$$

where A^* is not computable. Since $\Phi_A^{-1}(A^*) = \{A\}$, by Theorem 2.21,

$$A \in \text{MLR}^B \Leftrightarrow A^* \in \text{MLR}_\mu^B$$

for every $B \in 2^\omega$. Then there are exactly two $LR(\mu)$ -degrees:

$$\begin{aligned} \mathbf{0} &= \{B : A \in \text{MLR}^B\}, \\ \mathbf{1} &= \{B : A \notin \text{MLR}^B\}. \end{aligned}$$

Example 3. Let $A \oplus B \in \text{MLR} \cap \Delta_2^0$, and let Φ_A and Φ_B be the tally functionals for A and B , and let μ_0 and μ_1 be the measures induced by Φ_A and Φ_B , respectively. By Theorem 3.2 (ii),

$$\text{MLR}_{\mu_0} = \{\Phi_A(A)\} \cup \text{Atoms}_{\mu_0} \text{ and } \text{MLR}_{\mu_1} = \{\Phi_B(B)\} \cup \text{Atoms}_{\mu_1}$$

If we set $\nu := \frac{\mu_0 + \mu_1}{2}$, by Lemma 2.7, we have

$$\text{MLR}_\nu = \text{MLR}_{\mu_0} \cup \text{MLR}_{\mu_1} = \{\Phi_A(A), \Phi_B(B)\} \cup \text{Atoms}_{\mu_0} \cup \text{Atoms}_{\mu_1}.$$

Clearly, ν is trivial. There are exactly four $LR(\nu)$ -degrees; namely $\mathbf{0}$, \mathbf{a} , \mathbf{b} , and $\mathbf{1}$, where (using Theorem 2.21 and van Lambalgen's theorem)

$$\begin{aligned} \mathbf{0} &= \{X : A \in \text{MLR}^X \ \& \ B \in \text{MLR}^X\}, \\ \mathbf{a} &= \{X : A \in \text{MLR}^X \ \& \ B \notin \text{MLR}^X\}, \\ \mathbf{b} &= \{X : A \notin \text{MLR}^X \ \& \ B \in \text{MLR}^X\}, \\ \mathbf{1} &= \{X : A \notin \text{MLR}^X \ \& \ B \notin \text{MLR}^X\}. \end{aligned}$$

In particular, we have $\mathbf{0} < \mathbf{a} < \mathbf{1}$ and $\mathbf{0} < \mathbf{b} < \mathbf{1}$, but \mathbf{a} and \mathbf{b} are incomparable. Thus, $\mathcal{D}_{LR(\nu)}$ is isomorphic to the finite Boolean algebra on two atoms, pictured in Figure 1.




FIGURE 1. The finite Boolean algebra on two atoms

In the previous three examples we have defined trivial measures μ such that the associated $LR(\mu)$ -degrees are isomorphic to the finite Boolean algebra of one, two, and four elements, respectively. Thus, it is natural to consider whether there is such a measure for every finite Boolean algebra.

Theorem 5.1. *For every finite Boolean algebra (B, \leq) , there is a trivial measure $\mu \in \mathcal{M}_c$ such that*

$$(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)}) \cong (B, \leq).$$

Proof. First, some notation. Let $deg_{LR(\mu)}(X) = \{Y : X \equiv_{LR} Y\}$, and set

$$deg_{LR(\mu)}(X) \leq deg_{LR(\mu)}(Y)$$

if and only $X \leq_{LR} Y$. Now, we proceed in four steps:

Step 1: If n is the number of atoms of B , choose $A_1, A_2, \dots, A_n \in \text{MLR} \cap \Delta_2^0$ such that for each $J \subseteq \{1, \dots, n\}$, if

$$X_J = \bigoplus_{j \in J} A_j$$

then

$$A_i \in \text{MLR}^{X_J}$$

for every $i \notin J$. (This can be accomplished using Theorem 2.10.)

Step 2: For each A_i , let Φ_{A_i} be the tally functional defined in terms of the Δ_2^0 approximation of A_i , and define μ_i to be the measure induced by the tally functional Φ_{A_i} . Let $A_i^* = \Phi_{A_i}(A_i)$.

Step 3: Define $\mu := \frac{1}{n} \sum_{i=1}^n \mu_i$. It follows from Lemma 2.7 that

$$\text{MLR}_\mu = \bigcup_{i=1}^n \text{MLR}_{\mu_i} = \{A_1^*, A_2^*, \dots, A_n^*\} \cup \bigcup_{i=1}^n \text{Atoms}_{\mu_i}.$$

Step 4: We verify that for $J, K \subseteq \{1, \dots, n\}$, $deg_{LR(\mu)}(X_J) \leq deg_{LR(\mu)}(X_K)$ if and only if $J \subseteq K$. First, note that

$$\text{MLR}_\mu^{X_J} = \{A_i^* : i \notin J\} \ \& \ \text{MLR}_\mu^{X_K} = \{A_i^* : i \notin K\}.$$

Then

$$\begin{aligned} deg_{LR(\mu)}(X_J) \leq deg_{LR(\mu)}(X_K) &\Leftrightarrow \text{MLR}_\mu^{X_K} \subseteq \text{MLR}_\mu^{X_J} \\ &\Leftrightarrow \{A_i^* : i \notin K\} \subseteq \{A_i^* : i \notin J\} \\ &\Leftrightarrow J \subseteq K. \end{aligned}$$

Thus, $\mathcal{D}_{LR(\mu)} = \{deg_{LR(\mu)}(X_J) : J \subseteq \{1, \dots, n\}\}$ is isomorphic to the powerset of $\{1, \dots, n\}$, which is isomorphic to \mathcal{B} . \square

Theorem 5.1 can be improved further:

Theorem 5.2. *For every finite distributive lattice (L, \leq) , there is a trivial measure $\mu \in \mathcal{M}_c$ such that*

$$(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)}) \cong (L, \leq).$$

The following terminology will be useful in the proof of Theorem 5.2. Let L be a finite distributive lattice of n elements. We will consider L in terms of levels, where Level 1 consists of 1_L , Level 2 consists of the immediate predecessors of 1_L , Level 3 consists of the immediate predecessors of elements of Level 2, and so on. Since L has size n , there are only finitely

many levels (in fact, at most n levels), and since it is a lattice, the lowest level consists solely of $\mathbf{0}_L$.

Recall that for $a, b \in L$, the meet of a and b , denoted $a \wedge b$, is the greatest element in L such that $a \geq a \wedge b$ and $b \geq a \wedge b$. The element $c \in L$ is meet-reducible if there are $a, b > c$ such that $a \wedge b = c$, and it is meet-irreducible if it is not meet-reducible.

To prove Theorem 5.2, the idea is (i) construct a lattice of sets isomorphic to L , (ii) use these sets to define a collection of tally functionals, and (iii) define a measure in terms of these tally functionals, which will give rise to an LR -structure that is isomorphic to (L, \leq) . Let us first consider an example.

Let (L, \leq) be the finite distributive lattice given below in Figure 2.




FIGURE 2. The finite distributive lattice (L, \leq)

Now let $A \in \text{MLR} \cap \Delta_2^0$, and let $\{A_i\}_{i \in \omega}$ be such that

$$A = \bigoplus_{i \in \omega} A_i,$$

so that each $A_i \in \text{MLR} \cap \Delta_2^0$. Hereafter, the sequences A_1, A_2, \dots will be referred to as *basic sequences*. An important feature of these basic sequences is that each A_i is Martin-Löf random relative to a finite join of any basic sequences that differ from A_i (see Theorem 2.10).

We proceed by associating to each element at each level of L a set consisting of some of the A_i 's or joins of the A_i 's, yielding a finite distributive lattice of sets that is isomorphic to L , as in Figure 3.




FIGURE 3. A finite distributive lattice of sets isomorphic to \mathcal{L}

Level 1: We associate to the top element $1_{\mathcal{L}}$ the empty set.

Level 2: There are two elements in Level 2, and so we associate to one the set $\{A_1\}$ and to the other $\{A_2\}$.

Level 3: There are two elements in Level 3, one of which is meet-reducible and the other meet-irreducible. To the meet-reducible element, we associate the set $\{A_1, A_2\}$, and to the meet-irreducible element (which is below the element associated to the set $\{A_1\}$), we associate the set $\{A_1, A_1 \oplus A_3\}$, where A_3 is the first basic sequence in $\{A_i\}_{i \in \omega}$ (in the order given by the indices) that has not appeared in the construction thus far. Note that any sequence that derandomizes A_1 also derandomizes $A_1 \oplus A_3$, but not every element that derandomizes $A_1 \oplus A_3$ also derandomizes A_1 (such as A_3 itself).

Level 4: The only element at Level 4 is $0_{\mathcal{L}}$, the meet of the two Level 3 elements, and thus we associate to this element the set $\{A_1, A_2, A_1 \oplus A_3\}$.

Next, for each element in the set associated with $0_{\mathcal{L}}$, namely A_1 , A_2 , and $A_1 \oplus A_3$, let μ_{A_1} , μ_{A_2} , and $\mu_{A_1 \oplus A_3}$ be the measures induced by the tally functionals Φ_{A_1} , Φ_{A_2} , and $\Phi_{A_1 \oplus A_3}$ defined in terms of the Δ_2^0 approximations of A_1 , A_2 , and $A_1 \oplus A_3$. We define

$$\mu := \frac{1}{3}(\mu_{A_1} + \mu_{A_2} + \mu_{A_1 \oplus A_3}),$$

Let

$$\Phi_{A_1}(A_1) = A_1^*, \Phi_{A_2}(A_2) = A_2^*, \text{ and } \Phi_{A_1 \oplus A_3}(A_1 \oplus A_3) = (A_1 \oplus A_3)^*.$$

Then for any $X \in 2^\omega$,

$$\text{MLR}_\mu^X = \text{Atoms}_\mu \cup \mathcal{S},$$

where \mathcal{S} is equal to one of the following:

$$\emptyset, \{A_1^*\}, \{A_2^*\}, \{A_1^*, A_2^*\}, \{A_1^*, (A_1 \oplus A_3)^*\}, \text{ or } \{A_1^*, A_2^*, (A_1 \oplus A_3)^*\}.$$

Thus we have a one-to-one correspondence (that preserves \subseteq) between the above sets and those sets associated to the elements of L , and thus $(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)})$ is isomorphic to L . Now we proceed in full generality.

Proof of Theorem 5.2. Let L be a finite distributive lattice. We proceed as in the example. We first associate basic sequences and joins of basic sequences to elements of the various levels of L .

Level 1: We associate to the top element 1_L the empty set.

Level 2: To each of the $j \leq K$ elements in Level 2, we associate a singleton consisting of a basic sequence A_1, A_2, \dots, A_k .

Level $n + 1$: The set we associate to a Level $n + 1$ element depends on whether it is meet-reducible or meet-irreducible.

- The meet-reducible case: Let $a = b \wedge c$, where b and c are Level n elements. If \mathcal{S}_b is the set of sequences associated to b and \mathcal{S}_c is the set of sequences associated to c , then we associate the set $\mathcal{S}_b \cup \mathcal{S}_c$ to a . (Note: We will have to verify that this is well-defined, for it may be the case that there are Level n elements b' and c' that differ from b and c but also satisfy $b' \wedge c' = a$.)

- The meet-irreducible case: If a is meet-irreducible, then there is only one Level n element such that $a \leq b$. If \mathcal{S}_b is the set associated to b , then we proceed as follows. First, let $\{A_{i_1}, A_{i_2}, \dots, A_{i_k}\}$ be the collection of basic sequences appearing in S_b . That is, these are either elements of \mathcal{S}_b or are contained in joins in \mathcal{S}_b (so that, for instance, the basic sequences appearing in $\{A_1, A_2 \oplus A_3\}$ are A_1, A_2 , and A_3). Let $N \in \omega$ be the least such that the basic sequence A_N has not appeared in any set associated to an element of L . Then to a we associate the set

$$\left\{ \bigoplus_{j=1}^k A_{i_j} \oplus A_N \right\} \cup \mathcal{S}_b.$$

To verify that $\mathcal{S}_L = (\{\mathcal{S}_a : a \in L\}, \leq)$ is a finite distributive lattice isomorphic to (L, \leq) (where $\mathcal{S}_a \leq \mathcal{S}_b$ if and only if $\mathcal{S}_a \supseteq \mathcal{S}_b$), we first show that meets in \mathcal{S}_L are well-defined. First, suppose that $a, b, c, d \in L$ are distinct elements such $a = b \wedge c = c \wedge d$, as in Figure 4.




FIGURE 4. The case in which $a = b \wedge c = c \wedge d$

We claim that $b \vee c \neq c \vee d$. For otherwise, the lattice M_3 (pictured in Figure 5 below) would be embeddable into L , contradicting the fact that L is distributive.




FIGURE 5. The lattice M_3

Thus, we are in the situation as depicted by Figure 6.

We must also have $b \vee d \neq b \vee c$ and $b \vee d \neq c \vee d$, for otherwise M_3 is embeddable into L (for instance, Figure 7 shows the case that $b \vee d = c \vee d$).

If $e = b \vee c, f = c \vee d$, and $g = b \vee d$, then $b = e \wedge g, c = e \wedge f$, and $d = f \wedge g$, and thus none of b, c , or d is meet-irreducible.

Now, suppose that

$$(3) \quad \mathcal{S}_b \cup \mathcal{S}_c \neq \mathcal{S}_c \cup \mathcal{S}_d$$

Since $b = e \wedge g, c = e \wedge f$, and $d = f \wedge g$, the following hold:




FIGURE 6. $b \vee c \neq c \vee d$




FIGURE 7. The case that $b \vee d = c \vee d$

$$\begin{aligned}\mathcal{S}_b &= \mathcal{S}_e \cup \mathcal{S}_g \\ \mathcal{S}_c &= \mathcal{S}_e \cup \mathcal{S}_f \\ \mathcal{S}_d &= \mathcal{S}_f \cup \mathcal{S}_g.\end{aligned}$$

By (3), we have

$$\mathcal{S}_e \cup \mathcal{S}_f \cup \mathcal{S}_g = \mathcal{S}_b \cup \mathcal{S}_c \neq \mathcal{S}_c \cup \mathcal{S}_d = \mathcal{S}_e \cup \mathcal{S}_f \cup \mathcal{S}_g,$$

which is impossible. In the case where there are distinct Level n elements b, c, b', c' such that $a = b \wedge c = b' \wedge c'$, it follows that $a = b \wedge c'$, and so we apply the above argument to b, c, c' , and then to b, c, b' to conclude that

$$\mathcal{S}_b \cup \mathcal{S}_c = \mathcal{S}_{b'} \cup \mathcal{S}_{c'}.$$

Thus, meets are well-defined.

Next we show that the isomorphism between $\mathcal{S}_L = (\{\mathcal{S}_a : a \in L\}, \leq)$ and (L, \leq) holds level by level. In particular, we show that meets and joins are preserved level by level. First, it is clear that the top two levels of \mathcal{S}_L and L are isomorphic. Now suppose that \mathcal{S}_L and L are isomorphic from Level 1 to Level n . Having associated to Level n elements a and b the sets \mathcal{S}_a and \mathcal{S}_b , we associate the set $\mathcal{S}_a \cup \mathcal{S}_b$ to $a \wedge b$.

Suppose Level $n + 1$ elements a and b are associated with \mathcal{S}_a and \mathcal{S}_b . To show that $\mathcal{S}_{a \vee b}$, the set associated to $a \vee b$, is $\mathcal{S}_a \cap \mathcal{S}_b$, we consider three cases.

Case 1: First, if both a and b are meet-irreducible, then either (i) there is some Level n element c such that $a, b \leq c$, or (ii) there are distinct Level n elements c and d such that $a \leq c$ and $b \leq d$.

Subcase 1(i): By the procedure given above,

$$\mathcal{S}_a = \mathcal{S}_c \cup \{B \oplus A_\ell\}$$

and

$$\mathcal{S}_b = \mathcal{S}_c \cup \{B \oplus A_{\ell'}\},$$

where B is the join of the basic sequences appearing in \mathcal{S}_c and $A_\ell, A_{\ell'}$ are distinct basic sequences not contained in any set associated to elements of Levels $k \leq n$. Thus $\mathcal{S}_{a \vee b} = \mathcal{S}_c = \mathcal{S}_a \cap \mathcal{S}_b$.

Subcase 1(ii). In this subcase,

$$\mathcal{S}_a = \mathcal{S}_c \cup \{B_0 \oplus A_\ell\},$$

and

$$\mathcal{S}_b = \mathcal{S}_d \cup \{B_1 \oplus A_{\ell'}\},$$

where B_0 and B_1 are the joins of the basic sequences appearing in \mathcal{S}_c and \mathcal{S}_d , respectively, and $A_\ell, A_{\ell'}$ are distinct basic sequences not contained in any set associated to elements of Level k for any $k \leq n$. By induction, there is some $e \in L$ such that $e = c \vee d$ and $\mathcal{S}_e = \mathcal{S}_c \cap \mathcal{S}_d$. Then we have $e = a \vee b$ and

$$\mathcal{S}_a \cap \mathcal{S}_b = (\mathcal{S}_c \cup \{B_0 \oplus A_\ell\}) \cap (\mathcal{S}_d \cup \{B_1 \oplus A_{\ell'}\}) = \mathcal{S}_c \cap \mathcal{S}_d = \mathcal{S}_e = \mathcal{S}_{a \vee b}.$$

Case 2: If a is meet-irreducible but b is meet-reducible, then again there are two subcases to consider: Either (i) there is some Level n element c such that $a, b \leq c$, or (ii) there are distinct Level n elements c, d, e such that $a \leq c$ and $b = d \wedge e$.

Subcase 2(ii): We have

$$\mathcal{S}_a = \mathcal{S}_c \cup \{B \oplus A_\ell\},$$

where B is the join of the basic sequences appearing in \mathcal{S}_c and A_ℓ is a basic sequence not contained in any set associated to any element of Level k for any $k \leq n$, and

$$\mathcal{S}_b = \mathcal{S}_c \cup \mathcal{S}_d$$

for some Level n element $d \neq c$. Again it follows that

$$\mathcal{S}_a \cap \mathcal{S}_b = (\mathcal{S}_c \cup \{B \oplus A_\ell\}) \cap (\mathcal{S}_c \cup \mathcal{S}_d) = \mathcal{S}_c = \mathcal{S}_{a \vee b}.$$

Subcase 2(i): In this subcase, $c \vee (d \wedge e) = a \vee b$. As above,

$$\mathcal{S}_a = \mathcal{S}_c \cup \{B \oplus A_\ell\}$$

and

$$\mathcal{S}_b = \mathcal{S}_d \cup \mathcal{S}_e.$$

By the inductive hypothesis, we have $\mathcal{S}_c \cap (\mathcal{S}_d \cup \mathcal{S}_e) = \mathcal{S}_{c \vee (d \wedge e)}$, and thus

$$\mathcal{S}_a \cap \mathcal{S}_b = (\mathcal{S}_c \cup \{B \oplus A_\ell\}) \cap (\mathcal{S}_d \cup \mathcal{S}_e) = \mathcal{S}_c \cap (\mathcal{S}_d \cup \mathcal{S}_e) = \mathcal{S}_{c \vee (d \wedge e)} = \mathcal{S}_{a \vee b}.$$

Case 3: Lastly, in the case that a and b are both meet-reducible, either (i) there are distinct Level n elements c, d, e such $a = c \wedge d$, and $b = d \wedge e$ or (ii) there are distinct Level n elements c, d, e, f such that $a = c \wedge d$ and $b = e \wedge f$.

Subcase 3(i): Since $a, b \leq d$, it follows that $\mathcal{S}_a = \mathcal{S}_c \cup \mathcal{S}_d$, $\mathcal{S}_b = \mathcal{S}_d \cup \mathcal{S}_e$, $\mathcal{S}_c \cap \mathcal{S}_e = \emptyset$, and thus

$$\mathcal{S}_a \cap \mathcal{S}_b = (\mathcal{S}_c \cup \mathcal{S}_d) \cap (\mathcal{S}_d \cup \mathcal{S}_e) = \mathcal{S}_d = \mathcal{S}_{a \vee b}.$$

Subcase 3(ii): Note that $a \vee b = (c \wedge d) \vee (e \wedge f)$. Since $\mathcal{S}_a = \mathcal{S}_c \cup \mathcal{S}_d$ and $\mathcal{S}_b = \mathcal{S}_e \cup \mathcal{S}_f$, by the inductive hypothesis, it follows that

$$\mathcal{S}_a \cap \mathcal{S}_b = (\mathcal{S}_c \cup \mathcal{S}_d) \cap (\mathcal{S}_e \cup \mathcal{S}_f) = \mathcal{S}_{(c \vee d) \wedge (e \vee f)} = \mathcal{S}_{a \vee b}.$$

Having verified that \mathcal{S}_L is a finite distributive lattice, we now turn to defining the trivial measure μ . Let

$$\{B_1, \dots, B_k\}$$

be the set in \mathcal{S}_L associated to 0_L . By our construction, each B_i is either a basic sequence or the join of some basic sequences. Further, since the basic sequences are all in $\text{MLR} \cap \Delta_2^0$, and further, each is Martin-Löf random relative to the finite join of any number of basic sequences that differ from it, it follows from van Lambalgen's Theorem that each $B_i \in \text{MLR} \cap \Delta_2^0$.

Let Φ_i be the tally functional defined in terms of the Δ_2^0 approximation of B_i , and let $B_i^* = \Phi_i(B_i)$, so that $\text{MLR}_{\mu_i} = \{B_i^*\} \cup \text{Atoms}_{\mu_i}$. Further, given \mathcal{S} , one of the sets of sequences that is associated to some element of L , we define \mathcal{S}^* such that

$$B \in \mathcal{S} \text{ if and only if } B^* \in \mathcal{S}^*.$$

Setting

$$\mu := \frac{1}{k} \sum_{i=1}^k \mu_i,$$

we claim that $(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)}) \cong (\mathcal{S}_L, \leq)$. First we show that for each \mathcal{S}_a^* , there is some $X \in 2^\omega$ such that

$$\mathcal{S}_a^* \cup \text{Atoms}_\mu = \text{MLR}_\mu^X.$$

If we let $\text{RScope}(B) = \{X \in 2^\omega : B \in \text{MLR}^X\}$ be the *randomness scope* of B , note that by Theorem 2.21,

$$B_i \in \text{MLR}^X \Leftrightarrow B_i^* \in \text{MLR}_\mu^X,$$

and hence $X \in \text{RScope}(B_i)$ if and only if $B_i^* \in \text{MLR}_\mu^X$. Observe that $\lambda(\text{RScope}(B)) = 1$ for every $B \in \text{MLR}$, since by van Lambalgen's Theorem, $\text{MLR}^B \subseteq \text{RScope}(B)$ and $\lambda(\text{MLR}^B) = 1$.

If

$$\mathcal{S}_a^* = \{B_{i_1}^*, \dots, B_{i_j}^*\},$$

then

$$\mathcal{X} = \bigcap_{j=1}^k \text{RScope}(B_{i_j}) \neq \emptyset,$$

since the finite intersection of measure one sets has measure one. Thus for any $X \in \mathcal{X}$, we have $\text{MLR}_\mu^X = \mathcal{S}_a^* \cup \text{Atoms}_\mu$.

We claim that for each $X \in 2^\omega$, there is some $a \in L$ such that $\text{MLR}_\mu^X = \mathcal{S}_a^* \cup \text{Atoms}_\mu$. Let $\{A_1, \dots, A_k\}$ be the collection of basic sequences appearing in the elements of \mathcal{S}_L . For each $i \leq k$, let $a_i \in L$ be the element such that the basic sequence A_i first appears in \mathcal{S}_{a_i} . Further, for $j \leq k$, let $\{A_1, \dots, A_j\}$ be the basic sequences that make up the singletons assigned to Level 2 elements of L (which we'll call the *Level Two basic sequences*), and let $\{A_{j+1}, \dots, A_k\}$ be the basic sequences that added when we assign sets to meet-irreducible

elements of L (which we'll call the *meet-irreducible basic sequences*). For each $X \in 2^\omega$, there is some $J \subseteq \{1, \dots, j\}$ such that

$$X \in \bigcap_{i \in J} \text{RScope}(A_i) \ \& \ X \notin \bigcap_{i \in \{1, \dots, j\} \setminus J} \text{RScope}(A_i).$$

That is, J picks out the indices of the Level Two basic sequences that X fails to derandomize. Now if $J = \emptyset$, then as every $B_i \in \mathcal{S}_L$ is either a Level Two basic sequence or is the join of a Level Two basic sequence with some other sequence, it follows that $\text{MLR}_\mu^X = \text{Atoms}_\mu$. If $J \neq \emptyset$, then it follows from the construction that

$$\{A_i^* : i \in J\} \cup \text{Atoms}_\mu = \mathcal{S}_{\bigwedge_{i \in J} a_i}^* \cup \text{Atoms}_\mu \subseteq \text{MLR}_\mu^X.$$

Turning to the meet-irreducible basic sequences, there is some $K \subseteq \{j+1, \dots, k\}$ such that

$$X \in \bigcap_{i \in K} \text{RScope}(A_i) \ \& \ X \notin \bigcap_{i \in \{j+1, \dots, k\} \setminus K} \text{RScope}(A_i).$$

Now it may be that in the course of the construction, some A_i with $i \in K$ is joined to some A_ℓ with $\ell \notin J$ (or joined to some sequence with A_ℓ as a subsequence). This occurs when we associate a collection of sequences to a meet-irreducible element of L that is below the element of L to which we associated the Level Two basic sequence A_ℓ . Let

$$\widehat{K} = K \setminus \{i \in K : \mathcal{S}_{a_i} \supseteq \mathcal{S}_{a_j} \text{ for some } \ell \notin J\}.$$

Then if $\widehat{K} = \emptyset$, then $\text{MLR}_\mu^X = \mathcal{S}_{\bigwedge_{i \in J} a_i}^* \cup \text{Atoms}_\mu$. Otherwise, setting

$$a = \bigwedge_{i \in J} a_i \wedge \bigwedge_{i \in \widehat{K}} a_i,$$

it follows from the construction that $\text{MLR}_\mu^X = \mathcal{S}_a^* \cup \text{Atoms}_\mu$.

Thus, every \mathcal{S}_a^* is the collection of non-atoms in MLR_μ^X for some $X \in 2^\omega$, and for every $X \in 2^\omega$, there is some \mathcal{S}_a^* such that $\text{MLR}_\mu^X = \mathcal{S}_a^* \cup \text{Atoms}_\mu$. Since $X \leq_{LR(\mu)} Y$ if and only if $\text{MLR}_\mu^Y \subseteq \text{MLR}_\mu^X$, it follows that

$$(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)}) \cong (\mathcal{S}_L, \leq).$$

□

We conclude with several questions.

Question 5.3. *If $\mathcal{L} = (L, \leq)$ is an infinite, computable, distributive lattice, is there a trivial measure $\mu \in \mathcal{M}_c$ such that*

$$(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)}) \cong (L, \leq)?$$

Question 5.4. *Is there an example of a finite non-distributive lattice (L, \leq) and a trivial measure $\mu \in \mathcal{M}_c$ such that*

$$(\mathcal{D}_{LR(\mu)}, \leq_{LR(\mu)}) \cong (L, \leq)?$$

REFERENCES

- [BP12] Laurent Bienvenu and Christopher Porter. Strong reductions in effective randomness. *Theoret. Comput. Sci.*, 459:55–68, 2012. [7](#), [9](#)
- [DH10] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York, 2010. [2](#), [6](#), [7](#)
- [DNWY06] Rod Downey, Andre Nies, Rebecca Weber, and Liang Yu. Lowness and Π_2^0 nullsets. *J. Symbolic Logic*, 71(3):1044–1052, 2006. [6](#)
- [Kau91] Steven Kautz. *Degrees of Random Sets*. PhD thesis, Cornell University, 1991. [2](#), [5](#)
- [ML66] Per Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–619, 1966. [1](#)
- [Nie05] André Nies. Lowness properties and randomness. *Adv. Math.*, 197(1):274–305, 2005. [13](#)
- [Nie09] André Nies. *Computability and randomness*. Oxford Logic Guides. Oxford University Press, 2009. [2](#), [8](#)
- [NST05] André Nies, Frank Stephan, and Sebastiaan A. Terwijn. Randomness, relativization and Turing degrees. *J. Symbolic Logic*, 70(2):515–535, 2005. [5](#)
- [Sch69] Claus-Peter Schnorr. Eine Bemerkung zum Begriff der zufälligen Folge. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 14:27–35, 1969.
- [Sch71] Claus-Peter Schnorr. *Zufälligkeit und Wahrscheinlichkeit. Eine algorithmische Begründung der Wahrscheinlichkeitstheorie*. Lecture Notes in Mathematics, Vol. 218. Springer-Verlag, Berlin, 1971. [1](#)
- [Sch77] C.-P. Schnorr. A survey of the theory of random sequences. In *Basic problems in methodology and linguistics (Proc. Fifth Internat. Congr. Logic, Methodology and Philos. of Sci., Part III, Univ. Western Ontario, London, Ont., 1975)*, pages 193–211. Univ. Western Ontario Ser. Philos. Sci., Vol. 11. Reidel, Dordrecht, 1977. [1](#)
- [Soa87] Robert I. Soare. *Recursively enumerable sets and degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1987. A study of computable functions and computably generated sets. [2](#)
- [vL90] Michiel van Lambalgen. The axiomatization of randomness. *J. Symbolic Logic*, 55(3):1143–1167, 1990. [5](#)
- [ZL70] A. K. Zvonkin and L. A. Levin. The complexity of finite objects and the basing of the concepts of information and randomness on the theory of algorithms. *Uspehi Mat. Nauk*, 25(6(156)):85–127, 1970. [1](#)